

IMPORTANT INFORMATION

Instruct patients to read entire package insert before using RHINOCORT® Allergy Spray for complete instructions on how to get a bottle ready for use (primed), how to use the spray bottle, and how to clean the spray nozzle.

STEPS TO GET A BOTTLE READY FOR USE (PRIMED)

Before first use, a new bottle must be primed. To prime RHINOCORT® Allergy Spray:

- 1. Shake bottle gently.
- 2. Remove protective cap.
- 3. Hold bottle firmly, as shown in Figure A.
- 4. Prime the pump by quickly and firmly pressing down on the white collar while holding the base of the bottle with your thumb for 8 sprays. The pump is now ready to use.
- 5. a. If not used for two consecutive days, reprime the pump until a fine mist appears.
 - b. If not used for more than 14 days, rinse the spray tip of the pump using the cleaning steps listed on reverse side.
 After cleaning, reprime until a fine spray appears.

Figure A

INSTRUCTIONS FOR USE

Follow these instructions for daily use of RHINOCORT® Allergy Spray:

- 1. Gently blow nose to clear nostrils.
- 2. Shake bottle gently and remove cap.
- Hold bottle firmly (See Figure A).
 Insert spray tip into nostril (tip should not reach far into nose).
 Close other nostril with a finger and lean head slightly forward so the spray will aim toward the back of nose (See Figure B).
 DO NOT spray towards nasal septum (wall between two nostrils) (See Figure C).
- 4. For each spray, press down quickly and firmly on white collar while holding the base of the bottle with your thumb. Breathe gently inward.
- 5. After spraying into your nostril, lean your head slightly backward for a few seconds.
- 6. If a second spray is needed in the same nostril, repeat steps 3 through 5.

 For complete dosing instructions see "Directions for Use" on other side.
- 7. Repeat steps 3 through 6, for the other nostril.
- Avoid blowing your nose for 15 MINUTES after using RHINOCORT® Allergy Spray.
- 9. Wipe spray tip with a clean tissue and replace outer cap. Store the bottle in an upright position.



Figure C Wrong Position



spray towards nasal septum (wall between two nostrils)



MILH RODESONIDE

DIRECTIONS FOR USE

- Before first use, you must get a new bottle ready (primed). See other side.
- If bottle is not used for two consecutive days, reprime the pump until a fine spray appears.
- If bottle is not used for more than two weeks, rinse the spray tip of the pump using the cleaning steps listed below. After cleaning, reprime the pump until a fine spray appears.

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children
12 years of age and older

- once daily, spray 2 times into each nostril while sniffing gently
- once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 6 TO UNDER 12 YEARS OF AGE

• the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year

children	6	to	under
12 years	0	f a	ge

- an adult should supervise use
- once daily, spray 1 time into each nostril while sniffing gently
- if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children under 6 years of age

do not use

OTHER IMPORTANT INFORMATION FOR USE

- do not use more than directed
- if you forget a dose, do **not** double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

HOW TO CLEAN YOUR RHINOCORT® ALLERGY SPRAY

Rinse cap and spray tip regularly. To do this:

- 1. Remove cap (see Figure D) and gently pull spray tip away from the bottle.
- Wash only cap and spray tip in warm water and rinse in cold tap water.Shake to remove excess water.
- Allow cap and spray tip to air dry completely before reassembling the nasal spray.
- 4. Press and release spray nozzle until a fine spray is produced, taking care not to spray in the face.
- 5. If the spray tip becomes blocked, it can be cleared by repeating steps 1 through 3. Do not unblock the nasal applicator with a pin or other sharp object.







