

Painful gas upsetting your baby?

Tame tummy troubles with
Infants' Mylicon[®] Gas Relief Drops

**Babies can cry a lot during their first few months.
Could painful gas be the reason?**



*Gas is caused by
air bubbles trapped
in the tummy.*

- **Infant gas is very common**
Babies often swallow air as they learn to eat and when they cry. Sometimes formulas or certain foods passed from mom's diet into her breast milk don't agree with them.
- **Some babies are just more sensitive**
Each baby is different. One baby may be easily upset by things that other babies don't mind—like a soiled diaper, a loud noise, or a gassy belly.



Try these gas prevention tips

- **Pay attention to proper feeding**
To limit air intake, make sure baby's lips are tight and far back on your breast or bottle nipple. Keep baby upright for at least 30 minutes after eating.
- **Burp baby often**
Try burping after every 2 or 3 ounces (from a bottle) or when you switch breasts.

Using Infants' Mylicon[®] Gas Relief Drops

Dosing Information

WEIGHT	AGE	DOSAGE
Under 24 lbs	Under 2 years	0.3 mL
Over 24 lbs	Over 2 years	0.6 mL

Active ingredient: simethicone 20 mg (in each 0.3 mL)

- Gluten-free, with no saccharin, alcohol, or artificial flavors
- Can be used at every feeding, up to 12 times a day*
- Contains just a single active ingredient (simethicone); not absorbed into the bloodstream

To learn more, go to Mylicon.com



*Use only as directed.



Scan
to buy
now! 

